

Lecture 36

Solution Instability for the Explicit Method

As we saw in experiments using `myheat.m`, the solution can become unbounded unless the time steps are small. In this lecture we consider why.

Writing the Difference Equations in Matrix Form

If we use the boundary conditions $u(0) = u(L) = 0$ then the explicit method of the previous section has the form

$$u_{i,j+1} = ru_{i-1,j} + (1 - 2r)u_{i,j} + ru_{i+1,j} \quad \text{for } 1 \leq i \leq m - 1 \quad \text{and} \quad 0 \leq j \leq n - 1,$$

where $u_{0,j} = 0$ and $u_{m,j} = 0$. This is equivalent to the matrix equation

$$\mathbf{u}_{j+1} = A\mathbf{u}_j, \tag{36.1}$$

where \mathbf{u}_j is the column vector $(u_{1,j}, u_{2,j}, \dots, u_{m,j})'$ representing the state at the j th time step and A is the matrix

$$A = \begin{pmatrix} 1 - 2r & r & 0 & \cdots & 0 \\ r & 1 - 2r & r & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & r & 1 - 2r & r \\ 0 & \cdots & 0 & r & 1 - 2r \end{pmatrix}. \tag{36.2}$$

Unfortunately, this matrix can have a property which is very bad in this context. Namely, it can cause exponential growth of error unless r is small. To see how this happens, suppose that \mathbf{U}_j is the vector of correct values of u at time step t_j and \mathbf{E}_j is the error of the approximation \mathbf{u}_j , then

$$\mathbf{u}_j = \mathbf{U}_j + \mathbf{E}_j.$$

From (36.1), the approximation at the next time step will be

$$\mathbf{u}_{j+1} = A\mathbf{U}_j + A\mathbf{E}_j,$$

and if we continue for k steps,

$$\mathbf{u}_{j+k} = A^k\mathbf{U}_j + A^k\mathbf{E}_j.$$

The problem with this is the term $A^k\mathbf{E}_j$. This term is exactly what we would do in the power method for finding the eigenvalue of A with the largest absolute value. If the matrix A has eigenvalues with absolute value greater than 1, then this term will grow exponentially. Figure 36.1 shows the largest absolute value of an eigenvalue of A as a function of the parameter r for various sizes of the matrix A . As you can see, for $r > 1/2$ the largest absolute eigenvalue grows rapidly for any m and quickly becomes greater than 1.




Figure 36.1: Maximum absolute eigenvalue as a function of r for the matrix A from the explicit method for the heat equation calculated for matrices A of sizes $m = 2 \dots 10$. Whenever the maximum absolute eigenvalue is greater than 1 the method is unstable, i.e. errors grow exponentially with each step. When using the explicit method $r < 1/2$ is a safe choice.

Consequences

Recall that $r = ck/h^2$. Since this must be less than $1/2$, we have

$$k < \frac{h^2}{2c}.$$

The first consequence is obvious: k must be relatively small. The second is that h cannot be too small. Since h^2 appears in the formula, making h small would force k to be extremely small! A third consequence is that we have a converse of this analysis. Suppose $r < .5$. Then all the eigenvalues will be less than one. Recall that the error terms satisfy

$$\mathbf{u}_{j+k} = A^k \mathbf{U}_j + A^k \mathbf{E}_j.$$

If all the eigenvalues of A are less than 1 in absolute value then $A^k \mathbf{E}_j$ grows smaller and smaller as k increases. This is really good. Rather than building up, the effect of any error diminishes as time passes! From this we arrive at the following principle: **If the explicit numerical solution for a parabolic equation does not blow up, then errors from previous steps fade away!**

Finally, we note that if we have non-zero boundary conditions then instead of equation (36.1) we have

$$\mathbf{u}_{j+1} = A\mathbf{u}_j + r\mathbf{b}_j, \quad (36.3)$$

where the first and last entries of \mathbf{b}_j contain the boundary conditions and all the other entries are zero. In this case the errors behave just as before, if $r > 1/2$ then the errors grow and if $r < 1/2$ the errors fade away.

We can write a function program `myexpmatrix` that produces the matrix A in (36.2), for given inputs m and r . Without using loops we can use the `diag` command to set up the matrix:

```

function A = myexpmatrix(m,r)
    % produces the matrix for the explicit method for a parabolic equation
    % Inputs: m -- the size of the matrix
    %         r -- the main parameter, ck/h^2
    % Output: A -- an m by m matrix
    u = (1-2*r)*ones(m,1); % make a vector for the main diagonal
    v = r*ones(m-1,1);     % make a vector for the upper and lower diagonals
    A = diag(u) + diag(v,1) + diag(v,-1); % assemble
end

```

Test this using $m = 6$ and $r = .4, .6$. Check the eigenvalues and eigenvectors of the resulting matrices:

```

>> A = myexpmatrix(6,.6)
>> [v e] = eig(A)

```

What is the “mode” represented by the eigenvector with the largest absolute eigenvalue? How is that reflected in the unstable solutions?

Exercises

- 36.1 Let $L = \pi$, $T = 20$, $f(x) = .1 \sin(x)$, $g_1(t) = 0$, $g_2(t) = 0$, $c = .5$, and $m = 20$, as used in the program `myheat.m`. What value of n corresponds to $r = 1/2$? Try different n in `myheat.m` to find precisely when the method works and when it fails. Is $r = 1/2$ the boundary between failure and success? Hand in a plot of the last success and the first failure. Include the values of n and r in each.
- 36.2 Write a well-commented MATLAB **script** program that produces the graph in Figure 36.1 for $m = 4$. Your program should:
- define r values from 0 to 1,
 - for each r
 - create the matrix A by calling `myexppmatrix`,
 - calculate the eigenvalues of A ,
 - find the max of the absolute values, and
 - plot these numbers versus r .